arXiv:cond-mat/9606192v1 25 Jun 1996

Exact calculation of thermodynamical quantities of the integrable $t-J$ model

G. Jüttner and A. Klümper
Universität zu Köln
Institut für Theoretische Physik
Zülpicher Str. 77
D-50937, Germany

(January, 1996)

The specific heat and the compressibility for the integrable $t-J$ model are calculated showing Luttinger liquid behavior for low temperatures. A Trotter-Suzuki mapping and the quantum transfer matrix approach are utilized. Using an algebraic Bethe ansatz this method permits the exact calculation of the free energy and related quantities. A set of just two non-linear integral equations determining these quantities is studied for various particle densities and temperatures. The structure of the specific heat is discussed in terms of the elementary charge as well as spin excitations.

75.10.Jm, 05.50+q

The $t-J$ model has been studied intensively in recent years because of its importance as one of the most fundamental systems of strongly correlated electrons and its possible relevance for purely electronic mechanisms for high-T_c superconductivity [1,2]. The model describes the nearest-neighbor hopping of electrons with spin-exchange interaction. The effect of a strong repulsive on-site Coulomb interaction is modeled by the restriction of the Hilbert space to states without doubly occupied lattice sites. The one-dimensional Hamiltonian reads

$$\begin{aligned} H = &-t\sum_{j,\sigma}\mathcal{P}(c^{\dagger}_{j,\sigma}c_{j+1,\sigma}+c^{\dagger}_{j+1,\sigma}c_{j,\sigma})\mathcal{P} \\ &+J\sum_{j}(S_jS_{j+1}-n_jn_{j+1}/4) \end{aligned} \qquad (1)$$

where the projector $\mathcal{P}=\prod_j(1-n_{j\uparrow}n_{j\downarrow})$ ensures that double occupancies of sites are forbidden. At the supersymmetric point $2t=J$ the system was shown to be integrable [3,4] by the well-known Bethe ansatz. The ground state and excitation spectrum were investigated by Bares et al. [5]. Critical exponents were calculated by finite-size scaling and conformal field theory [6,7].

According to the pioneering work of Takahashi [8] thermodynamical properties of general integrable systems are described by an infinite set of coupled integral equations reflecting the existence of infinitely many different rapidity patterns. For the $t-J$ model such integral equations were formulated in [9,10], but not solved. We like to point out two shortcomings of the traditional approach. First, the method consists of a rather direct evaluation of the partition function by taking into account all excited states of the Hamiltonian. The excitations are derived on the basis of the so-called string conjecture which describes the solutions of the Bethe ansatz equations. However, the validity of the string conjecture is quite controversial. Second, the calculation of the asymptotics of correlation functions is not possible within the traditional approach.

Our approach takes a different route following [11,12] to overcome the mentioned problems. We do not directly study the Hamiltonian at finite temperatures. Instead, we employ a convenient Trotter-Suzuki mapping [11] to an exactly solvable two-dimensional classical model. Furthermore, the corresponding thermodynamical quantities are expressed by eigenvalues of the so-called quantum transfer matrix. The largest eigenvalue, for example, directly yields the free energy. The next-largest eigenvalues provide the correlation lengths. This represents the important advantage compared with the traditional thermodynamical Bethe ansatz requiring all eigenvalues of the Hamiltonian. The new method using the Trotter-Suzuki mapping as well as the quantum transfer matrix approach has been applied to several quantum systems [11–16] notably the Hubbard model in [17].

The $t-J$ quantum chain belongs to a family of integrable fermionic systems which are related to the classical Perk-Schulz model [18], a multi-component generalization of the well-known six-vertex model. The detailed derivation of the quantum transfer matrix and the study of the corresponding eigenvalue problem by the quantum inverse scattering method will be described in a separate publication [19,20]. Thus, we confine ourselves to the main points.

The Perk-Schultz model is exactly solvable as the Boltzmann weights satisfy the Yang-Baxter equation. This in turn implies the commutation of all row-to-row transfer matrices for arbitrary spectral parameters u, v: $T(u)T(v)=T(v)T(u)$. Consequently, the Hamiltonian is integrable. We have

$$H=\frac{d}{du}\ln T(u)_{|u=0} \quad \text{with} \quad T(u)=T_R\,e^{uH+\mathcal{O}(u^2)}, \qquad (2)$$

where T_R is a right-shift operator. By means of the substitution $u=-\beta/N$ (β being the inverse temperature) we find

$$\left(T(-\beta/N)\overline{T}(-\beta/N)\right)^{N/2}=e^{-\beta H+\mathcal{O}(1/N)}, \qquad (3)$$

where $\overline{T}$ is the row-to-row transfer matrix of the Perk-Schultz model after a 90^o rotation. It satisfies (2) with

T, T_R replaced by $\overline{T}$, T_R^{-1}. The partition function of the quantum system

$$Z = \lim_{N\to\infty} \mathrm{Trace}\big(T(-\beta/N)\overline{T}(-\beta/N)\big)^{N/2} \tag{4}$$

is identical to the partition function of the inhomogeneous Perk-Schulz model with alternating rows [19]. The technically more convenient column-to-column transfer matrix of such a system is often referred to as the quantum transfer matrix (QTM). This matrix is shown to be embedded into a commuting family [19,20] of staggered transfer matrices given by alternating products of unrotated and rotated vertices. We still have the same intertwiner as in the case of the homogeneous lattice. In contrast to the work in [17] for the Hubbard model the integrability is preserved within the present approach. Thus, the eigenvalues of the QTM are analytic functions of the spectral parameter. This fact represents the important advantage leading to simple calculations which can be generalized to other models. The free energy can be calculated by the largest eigenvalue Λ of the QTM $f = -(\log \Lambda)/\beta$ and the next-leading eigenvalues yield the correlation lengths.

We treat the quantum transfer matrix by using an algebraic Bethe ansatz [19]. The eigenvalues $\Lambda(v)$ of $T(v)$ are shown to be analytic functions of the spectral parameter v with explicit representation

$$\Lambda(v) = \lambda_1(v) + \lambda_2(v) + \lambda_3(v), \tag{5}$$

where

$$\begin{aligned}
\lambda_1(v) &= \prod_j \frac{v - w_j + i}{v - w_j}\Big[(v - iu - i)(v + iu)\Big]^{N/2} e^{\beta\mu},\\
\lambda_2(v) &= \prod_k \frac{v - v_k - i}{v - v_k}\Big[(v + iu + i)(v - iu)\Big]^{N/2} e^{\beta\mu},\\
\lambda_3(v) &= \prod_j \frac{v - w_j + i}{v - w_j}\,(v + iu)^{N/2} \times\\
&\quad \prod_k \frac{v - v_k - i}{v - v_k}\,(v - iu)^{N/2}
\end{aligned}$$

and v_j and w_j are Bethe ansatz rapidities, μ denotes the chemical potential measured from the lower edge of the band. The defining relations for the rapidities, the so-called Bethe ansatz equations are equivalent to the analyticity of $\Lambda(v)$, i.e. the absence of poles. We use this analyticity to determine the largest eigenvalue by a finite set of non-linear integral equations [19,20]. This also allows for taking the limit $N \to \infty$ analytically. We define the following auxiliary functions (with arbitrary but fixed $0 < \gamma < 1$)

$$\begin{aligned}
b(x) &= \frac{\lambda_1(x + i\gamma)}{\lambda_2(x + i\gamma) + \lambda_3(x + i\gamma)},\\
c(x) &= \frac{\lambda_1(x)\lambda_2(x)}{\lambda_3(x)\big(\lambda_1(x) + \lambda_2(x) + \lambda_3(x)\big)}.
\end{aligned}$$

Using this definition and the analyticity property of the eigenvalue $\Lambda(v)$ a set of functional equations for b and c is derived. This set can be transformed into integral equations by standard applications of the Fourier transform. Such an approach should be applicable to all integrable models. After some lengthy calculations [20] we obtain the following relations for the two auxiliary functions

$$\begin{aligned}
\log b(x) &= -2\pi\beta\Psi_b(x + i\gamma) + \beta\mu -\\
&\qquad \Psi_b * \log(1 + \overline{b})|_{x+2i\gamma} - \Psi_b * \log(1 + c)|_{x+i\gamma},\\
\log c(x) &= -2\pi\beta\Psi_c(x) + 2\beta\mu -\\
&\qquad 2\Re\big(\Psi_b * \log(1 + \overline{b})|_{x+i\gamma}\big) - \Psi_c * \log(1 + c)|_x,
\end{aligned}$$

with

$$\Psi_b(x) = 1/\big(2\pi x(x - i)\big), \quad \Psi_c(x) = 2/\big(2\pi(x^2 + 1)\big),$$

where $*$ denotes the convolution $f * g = \int f(x-y)g(y)dy$ taken at the indicated arguments x, $x + i\gamma$ and $x + 2i\gamma$. The solution of the integral equations provides the largest eigenvalue via:

$$\log \Lambda = -\log c(0) + 2\beta\mu. \tag{6}$$

In order to calculate derivatives of the thermodynamical potential one can avoid numerical differentiations by utilizing similar integral equations [20] guaranteeing the same numerical accuracy as for of the free energy.

In the low-temperature and low-density limit (small μ and T/μ) we obtain an analytical expression for the grand canonical potential $f = -(\log \Lambda)/\beta$

$$f = -\int_{-\infty}^{\infty} \frac{dx}{\pi\beta^{3/2}}\,\log\big(1 + e^{-x^2 + \beta\mu}\big), \tag{7}$$

in agreement with the numerical results. Asymptotically we find

$$f = -\frac{4}{3\pi}\mu^{3/2} - \frac{\pi}{6\sqrt{\mu}}T^2 + o(\mu^{3/2}, T^2/\sqrt{\mu}), \tag{8}$$

implying a particle density $n = -\partial f/\partial\mu$ and entropy $S = -\partial f/\partial T$:

$$n = \frac{2}{\pi}\sqrt{\mu}, \quad S = \frac{2}{3n}T. \tag{9}$$

Moreover, due to predictions of Luttinger liquid theory and conformal field theory we expect for the low-temperature asymptotics

$$f = f_0 - \left(\frac{\pi c_s}{6 v_s} + \frac{\pi c_c}{6 v_c}\right)T^2 \tag{10}$$

where $v_{s,c}$ and $c_{s,c}$ are the velocities and central charges ($c_{s,c} = 1$) for the elementary spin and charge excitations. For small particle densities we have $v_{s,c} = \pi n$ [5], thus (8) and (10) are consistent.

On the other hand, the simple high-temperature limit for finite μ immediately leads to $S = \ln 3$ (and $n = 2/3$) as is expected by counting the degrees of freedom per lattice site. Therefore, the analytical benchmarks for $T \to 0$ and $T \to \infty$ imply the consistency of our approach.

Now we consider various thermodynamical quantities at intermediate temperatures by solving the integral equations numerically. Figure 1 shows the specific heat as a function of T for different fixed particle densities. First of all, we note a linear temperature dependence at low T. According to conformal field theory (10) the coefficient is given by $\pi(1/v_s + 1/v_c)/3$. Our numerical data are consistent with this expression. Furthermore, we observe two maxima with changing dominance for increasing particle density n. The nature of this structure can be understood from the elementary excitations of the system. In the groundstate the particles are bound in singlet pairs with binding energies varying from 0 to some density dependent value. There are two types of excitations. First, there are charge excitations due to energy-momentum transfer onto individual pairs. Second, there are excitations due to the breaking of pairs. The latter excitation is of spin type at lower excitation energies, but changes the character at higher (density dependent) energies to charge type as it describes the motion of single particles. Therefore, the first and second maximum at lower densities (figure 1.a) are caused by charge excitations due to pairs and single particles, respectively. At higher densities (figure 1.b) the maximum at lower temperatures is dominated by excitations of pairs whereas the second one at higher temperatures is caused by spin excitations. For increasing concentration the spin contribution becomes dominant as the charge excitations freeze out. This is in accordance with the limiting case $n = 1$ leading to the spin-1/2 Heisenberg chain. The missing spin structure in the specific heat at low and intermediate densities is found at quite low temperatures shown in figure 2.

It is worthwhile to compare these results with the findings for the Hubbard model investigated by the traditional thermodynamical Bethe ansatz [21]. The structure found in the specific heat is explained by spin and charge excitations which do not change their character in contrast to the $t - J$ model. For certain densities a low-temperature charge peak was found which is caused by single particle excitations. A charge peak at higher temperatures appears because of excitations due to doubly occupied lattice sites with energies of the order U for large Coulomb interaction.

To conclude our investigation we present numerical results for other thermodynamical quantities. Figure 3 presents the entropy and the compressibility $\kappa = \partial n/\partial\mu$ for various particle densities. Note the divergent low-temperature compressibility for particle densities $n \to 0$ and $n \to 1$.

In summary, we have derived eigenvalue equations for the quantum transfer matrix of the $t - J$ model. This approach permits the *exact* analytical as well as numerical calculation of thermodynamical quantities. Instead of solving an infinite set of integral equations – as is necessary in the traditional thermodynamical Bethe ansatz – we have to solve integral equations for only *two* functions. We have considered analytically certain low and high-temperature limits verifying the values predicted by conformal field theory. Moreover, the case of intermediate temperatures was treated numerically. As shown, the specific heat and compressibility display an interesting behavior in dependence on particle density and temperature.

The authors acknowledge financial support by the *Deutsche Forschungsgemeinschaft* under grant No. Kl 645/3-1.

---

[1] P. W. Anderson, Science **235**, 1196 (1987)

[2] F. C. Zhang and T. M. Rice, Phys. Rev. B **37**, 3759 (1988)

[3] B. Sutherland, Phys. Rev. B **12**, 3795 (1975)

[4] P. Schlottmann, Phys. Rev. B **36**, 5177 (1987)

[5] P. A. Bares, G. Blatter and M. Ogata Phys. Rev. B **44**, 130 (1991)

[6] N. Kawakami and S. K. Yang, Phys. Rev. Lett. **65**, 2309 (1990)

[7] R. Z. Bariev, A. Klümper, A. Schadschneider and J. Zittartz, Z. Phys. B **96**, 95 (1995)

[8] M. Takahashi, Prog. Theor. Phys. **46**, 401 (1971)

[9] E. D. Williams, *Thermodynamics and Excitations of the Supersymmetric $t - J$ Model*, Stony Brook preprint ITP-SB-92-72, e-print: cond-mat/9304009

[10] P. Schlottmann, J. Phys. C **4**, 7565 (1992)

[11] M. Suzuki, Phys. Rev. B. **31**, 2957 (1985)

[12] M. Suzuki and M. Inoue, Prog. Theor. Phys. **78**, 787 (1987)

[13] J. Suzuki, Y. Akutsu and M. Wadati, J. Phys. Soc. Japan **59**, 2667 (1990)

[14] M. Takahashi, Phys. Rev. B **43** (1991) 5788, Phys. Rev. B **44**, 12382 (1991)

[15] A. Klümper, Z. Phys. B **91** (1993) 507, Ann. Physik 1, 540 (1992)

[16] C. Destri and H. J. de Vega, Phys. Rev. Lett. **69**, 2313 (1992)

[17] A. Klümper and R. Z. Bariev, Nucl. Phys. B **458**, 623 (1996)

[18] C. L. Schulz, Phys. Rev. Lett. **46**, 629 (1981)

[19] A. Klümper, T. Wehner and J. Zittartz, in preparation: *Thermodynamics of the quantum Perk-Schulz model*

[20] G. Jüttner and A. Klümper, in preparation: *Exact thermodynamics of the integrable $t - J$ model*

[21] T. Usuki, N. Kawakami and A. Okiji, J. Phys. Soc. Japan **59**, 1357 (1990)



FIG. 1. Specific heat as function of T for different particle densities n with $n \leq 0.6$ and $n \geq 0.6$.



FIG. 2. Specific heat at low temperatures T for intermediate particle densities n.



FIG. 3. Entropy S and compressibility κ versus T for different densities n.